

JEE–MAIN EXAMINATION – JANUARY 2025

(HELD ON WEDNESDAY 22nd JANUARY 2025)

TIME : 9:00 AM TO 12:00 NOON

MATHEMATICS

TEST PAPER WITH SOLUTION

SECTION-A

1. The number of non-empty equivalence relations on the set $\{1,2,3\}$ is :

- (1) 6 (2) 7
- (3) 5 (4) 4

Ans. (3)

Sol. Let R be the required relation

$$A = \{(1, 1) (2, 2), (3, 3)\}$$

(i) $|R| = 3$, when $R = A$

(ii) $|R| = 5$, e.g. $R = A \cup \{(1, 2), (2, 1)\}$

Number of R can be [3]

(iii) $R = \{1, 2, 3\} \times \{1, 2, 3\}$

Ans. (5)

2. Let $f : \mathbf{R} \rightarrow \mathbf{R}$ be a twice differentiable function such that $f(x + y) = f(x) f(y)$ for all $x, y \in \mathbf{R}$. If $f'(0) = 4a$ and f satisfies $f''(x) - 3a f'(x) - f(x) = 0$, $a > 0$, then the area of the region

$R = \{(x,y) \mid 0 \leq y \leq f(ax), 0 \leq x \leq 2\}$ is :

- (1) $e^2 - 1$ (2) $e^4 + 1$
- (3) $e^4 - 1$ (4) $e^2 + 1$

Ans. (1)

Sol. $f(x + y) = f(x).f(y)$

$$\Rightarrow f(x) = e^{\lambda x} \quad f'(0) = 4a$$


$$\Rightarrow f'(x) = \lambda e^{\lambda x} \Rightarrow \lambda = 4a$$

$$\text{So, } f(x) = e^{4ax}$$

$$f''(x) - 3af'(x) - f(x) = 0$$

$$\Rightarrow \lambda^2 - 3a\lambda - 1 = 0$$

$$\Rightarrow 16a^2 - 12a^2 - 1 = 0 \Rightarrow 4a^2 = 1 \Rightarrow a = \frac{1}{2}$$



$$F(x) = e^{2x}$$

$$\text{Area} = \int_0^2 e^x dx = e^2 - 1$$

3. Let the triangle PQR be the image of the triangle with vertices (1,3), (3,1) and (2, 4) in the line $x + 2y = 2$. If the centroid of ΔPQR is the point (α, β) , then $15(\alpha - \beta)$ is equal to :

- (1) 24 (2) 19
- (3) 21 (4) 22

Ans. (4)

Sol. Let 'G' be the centroid of Δ formed by (1, 3) (3, 1) & (2, 4)

$$G \cong \left(2, \frac{8}{3}\right)$$

Image of G w.r.t. $x + 2y - 2 = 0$

$$\frac{\alpha - 2}{1} = \frac{\beta - \frac{8}{3}}{2} = -2 \frac{\left(2 + \frac{16}{3} - 2\right)}{1 + 4}$$

$$= -2 \left(\frac{16}{3}\right)$$

$$\Rightarrow \alpha = \frac{-32}{15} + 2 = \frac{-2}{15}, \quad \beta = \frac{-32 \times 2}{15} + \frac{8}{3} = \frac{-24}{15}$$

$$15(\alpha - \beta) = -2 + 24 = 22$$

4. Let z_1, z_2 and z_3 be three complex numbers on the circle

$|z| = 1$ with $\arg(z_1) = \frac{-\pi}{4}$, $\arg(z_2) = 0$ and $\arg(z_3) = \frac{\pi}{4}$.

If $|z_1 \bar{z}_2 + z_2 \bar{z}_3 + z_3 \bar{z}_1|^2 = \alpha + \beta \sqrt{2}$, $\alpha, \beta \in \mathbf{Z}$, then the value of $\alpha^2 + \beta^2$ is :

- (1) 24 (2) 41
- (3) 31 (4) 29

Ans. (4)

Sol. $Z_1 = e^{-i\pi/4}$, $Z_2 = 1$, $Z_3 = e^{i\pi/4}$

$$|z_1 \bar{z}_2 + z_2 \bar{z}_3 + z_3 \bar{z}_1|^2 = \left| e^{-i\pi/4} \times 1 + 1 \times e^{-i\pi/4} + e^{i\pi/4} \times e^{i\pi/4} \right|^2$$

$$= \left| e^{-i\pi/4} + e^{-i\pi/4} + e^{i\pi/4} \right|^2$$

$$= \left| 2e^{-i\pi/4} + i \right|^2 = \left| \sqrt{2} - \sqrt{2}i + i \right|^2$$

$$= (\sqrt{2})^2 + (1 - \sqrt{2})^2 = 2 + 1 + 2 - 2\sqrt{2} = 5 - 2\sqrt{2}$$

$$\alpha = 5, \beta = -2$$

$$\Rightarrow \alpha^2 + \beta^2 = 29$$

$$\Rightarrow (3\mu - 2\lambda + 1)2 + (4\mu - 3\lambda + 2)3 + (5\mu - 4\lambda + 2)4 = 0$$

$$38\mu - 29\lambda + 16 = 0 \quad \dots(1)$$

PQ \perp L₂

$$\Rightarrow (3\mu - 2\lambda + 1)3 + (4\mu - 3\lambda + 2)4 + (5\mu - 4\lambda + 2)5 = 0$$

$$50\mu - 38\lambda + 21 = 0 \quad \dots(2)$$

By (1) & (2)

$$\lambda = \frac{1}{3}; \quad \mu = \frac{-1}{6}$$

$$\therefore P\left(\frac{5}{3}, 3, \frac{13}{3}\right) \text{ \& } Q\left(\frac{3}{2}, \frac{10}{3}, \frac{25}{6}\right)$$

Line PQ

$$\frac{x - \frac{5}{3}}{\frac{1}{6}} = \frac{y - 3}{\frac{-1}{3}} = \frac{z - \frac{13}{3}}{\frac{1}{6}}$$

$$\frac{x - \frac{5}{3}}{1} = \frac{y - 3}{-2} = \frac{z - \frac{13}{3}}{1}$$

$$\text{Point}\left(\frac{14}{3}, -3, \frac{22}{3}\right)$$

lies on the line PQ

9. The product of all solutions of the equation

$$e^{5(\log_e x)^2 + 3} = x^8, \quad x > 0, \text{ is :}$$

$$(1) e^{8/5} \qquad (2) e^{6/5}$$

$$(3) e^2 \qquad (4) e$$

Ans. (1)

Sol. $e^{5(\ln x)^2 + 3} = x^8$

$$\Rightarrow \ln e^{5(\ln x)^2 + 3} = \ln x^8$$

$$\Rightarrow 5(\ln x)^2 + 3 = 8 \ln x$$

$$(\ln x = t)$$

$$\Rightarrow 5t^2 - 8t + 3 = 0$$

$$t_1 + t_2 = \frac{8}{5}$$

$$\ln x_1 x_2 = \frac{8}{5}$$

$$x_1 x_2 = e^{8/5}$$

10. If $\sum_{r=1}^n T_r = \frac{(2n-1)(2n+1)(2n+3)(2n+5)}{64}$, then

$\lim_{n \rightarrow \infty} \sum_{r=1}^n \left(\frac{1}{T_r}\right)$ is equal to :

(1) 1 (2) 0

(3) $\frac{2}{3}$ (4) $\frac{1}{3}$

Ans. (3)

Sol. $T_n = S_n - S_{n-1}$

$$\Rightarrow T_n = \frac{1}{8}(2n-1)(2n+1)(2n+3)$$

$$\Rightarrow \frac{1}{T_n} = \frac{8}{(2n-1)(2n+1)(2n+3)}$$

$$\lim_{n \rightarrow \infty} \sum_{r=1}^n \frac{1}{T_r} = \lim_{n \rightarrow \infty} 8 \sum_{r=1}^n \frac{1}{(2n-1)(2n+1)(2n+3)}$$

$$= \lim_{n \rightarrow \infty} \frac{8}{4} \sum \left(\frac{1}{(2n-1)(2n+1)} - \frac{1}{(2n+1)(2n+3)} \right)$$

$$= \lim_{n \rightarrow \infty} 2 \left[\left(\frac{1}{1.3} - \frac{1}{3.5} \right) + \left(\frac{1}{3.5} - \frac{1}{5.7} \right) + \dots \right]$$

$$= \frac{2}{3}$$

11. From all the English alphabets, five letters are chosen and are arranged in alphabetical order. The total number of ways, in which the middle letter is 'M', is :

(1) 14950 (2) 6084

(3) 4356 (4) 5148

Ans. (4)

Sol. $\underbrace{AB}_{12} \quad \underbrace{MN \dots \dots \dots Z}_{13}$


$$= \underbrace{{}^{12}C_2}_{\text{Selection of two letters before M}} \times \underbrace{{}^{13}C_2}_{\text{Selection of two letters after M}} = 5148$$

12. Let $x = x(y)$ be the solution of the differential equation $y^2 dx + \left(x - \frac{1}{y}\right) dy = 0$. If $x(1) = 1$, then

$x\left(\frac{1}{2}\right)$ is :

(1) $\frac{1}{2} + e$ (2) $\frac{3}{2} + e$

(3) $3 - e$ (4) $3 + e$



$$\text{Area} = \int_{-\sqrt{3}}^1 (-6x^2 - 2 + 20) dx + \int_1^2 (4 - 12x + 20) dx$$

$$16 + 12\sqrt{3} + 6 = 22 + 12\sqrt{3}$$

22. If $\sum_{r=0}^5 \frac{{}^{11}C_{2r+1}}{2r+2} = \frac{m}{n}$, $\gcd(m, n) = 1$, then $m - n$ is equal to _____.

Ans. (2035)

$$\text{Sol. } \int_0^1 (1+x)^{11} dx = \left[C_0 x + \frac{C_1 x^2}{2} + \frac{C_2 x^3}{3} + \dots \right]_0^1$$

$$\frac{2^{12} - 1}{12} = C_0 + \frac{C_1}{2} + \frac{C_2}{3} + \frac{C_3}{4} + \dots$$

$$\int_{-1}^0 (1+x)^{11} dx = \left[C_0 x + \frac{C_1 x^2}{2} + \frac{C_2 x^3}{3} + \dots \right]_{-1}^0$$

$$\frac{1}{12} = C_0 - \frac{C_1}{2} + \frac{C_2}{3} - \frac{C_3}{4} + \dots$$

$$\frac{2^{12} - 2}{12} = 2 \left(\frac{C_1}{2} + \frac{C_3}{4} + \frac{C_5}{6} + \dots \right)$$

$$\frac{C_1}{2} + \frac{C_3}{4} - \frac{C_5}{6} + \dots = \frac{2^{11} - 1}{12} = \frac{2047}{12}$$

23. Let A be a square matrix of order 3 such that $\det(A) = -2$ and $\det(3\text{adj}(-6\text{adj}(3A))) = 2^{m+n} \cdot 3^{mn}$, $m > n$. Then $4m + 2n$ is equal to _____.

Ans. (34)

$$\text{Sol. } |A| = -2$$

$$\det(3\text{adj}(-6\text{adj}(3A)))$$

$$= 3^3 \det(\text{adj}(-\text{adj}(3A)))$$

$$= 3^3 (-6)^6 (\det(3A))^4$$

$$= 3^{21} \times 2^{10}$$

$$m + n = 10$$

$$mn = 21$$

$$m = 7; n = 3$$

24. Let $L_1 : \frac{x-1}{3} = \frac{y-1}{-1} = \frac{z+1}{0}$ and

$$L_2 : \frac{x-2}{2} = \frac{y}{0} = \frac{z+4}{\alpha}, \alpha \in \mathbb{R},$$

be two lines, which intersect at the point B . If P is the foot of perpendicular from the point $A(1, 1, -1)$ on L_2 , then the value of $26 \alpha(PB)^2$ is _____.

Ans. (216)

Sol. Point B

$$(3\lambda + 1, -\lambda + 1, -1) \equiv (2\mu + 2, 0, \alpha\mu - 4)$$

$$3\lambda + 1 = 2\mu + 2$$

$$-\lambda + 1 = 0$$

$$-1 = \alpha\mu - 4$$

$$\lambda = 1, \mu = 1, \alpha = 3$$

$$B(4, 0, -1)$$

$$\text{Let Point 'P' is } (2\delta + 2, 0, 3\delta - 4)$$

$$\text{Dir's of } AP < 2\delta + 1, -1, 3\delta - 3 >$$

$$AP \perp L_2 \Rightarrow \delta = \frac{7}{13}$$

$$P\left(\frac{40}{13}, 0, \frac{-31}{13}\right)$$

$$2\sigma\delta(PB)^2 = 26 \times 3 \times \left(\frac{144}{169} + \frac{324}{169}\right)$$

$$= 216$$

25. Let \vec{c} be the projection vector of $\vec{b} = \lambda\hat{i} + 4\hat{k}$, $\lambda > 0$, on the vector $\vec{a} = \hat{i} + 2\hat{j} + 2\hat{k}$. If $|\vec{a} + \vec{c}| = 7$, then the area of the parallelogram formed by the vectors \vec{b} and \vec{c} is _____.

Ans. (16)

$$\text{Sol. } \vec{c} = \left(\frac{\vec{b} \cdot \vec{a}}{|\vec{a}|} \right) \frac{\vec{a}}{|\vec{a}|}$$

$$= \left(\frac{\lambda + 8}{9} \right) (\hat{i} + 2\hat{j} + 2\hat{k})$$

$$|\vec{a} + \vec{c}| = 7 \Rightarrow \lambda = 4$$

Area of parallelogram

$$= \left| \vec{b} \times \vec{c} \right| = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 4 & 8 & 8 \\ 4 & 0 & 4 \end{vmatrix}$$

$$= 16$$